

# BLANCHARD AND KAHN'S (1980) SOLUTION FOR A LINEAR RATIONAL EXPECTATIONS MODEL WITH ONE STATE VARIABLE AND ONE JUMP VARIABLE: THE CORRECT FORMULA\*

by

ROBERT KOLLMANN

*ECARES, Université Libre de Bruxelles, Université Paris-Est and CEPR*  
and

STEFAN ZEUGNER<sup>†</sup>

*DG-ECFIN, European Commission*

This note corrects Blanchard and Kahn's (1980) formula for the solution of a linear dynamic rational expectations model with one predetermined and one non-predetermined endogenous variable.

## 1 INTRODUCTION

In their classical paper, Blanchard and Kahn (1980) [BK] derived the solution for an important class of dynamic linear rational expectations models. The BK algorithm has become a standard tool for economic modelers.<sup>1</sup> In general, the model solution is analytically intractable. However, as shown by BK, a model with one predetermined and one non-predetermined endogenous variable can be handled analytically, which facilitates an intuitive understanding of the solution. That special case is important (and of pedagogical interest) as it includes, e.g., the basic RBC model with fixed labor. This note shows that the formula provided by BK, for this special case, includes an error; we also provide the correct formula.<sup>2</sup>

\* Manuscript received 17.9.16; final version received 15.11.16.

<sup>†</sup> R. Kollmann is also affiliated with the Globalization and Monetary Policy Institute (Federal Reserve Bank of Dallas) and with the Centre for Applied Macroeconomic Analysis (Australian National University). We thank Xavier Gabaix for encouraging us to circulate this note. The research leading to these results has received funding from the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement no. 612796, 'Integrated Macro-Financial Modelling for Robust Policy Design' (MACFINROBODS).

<sup>1</sup>The BK algorithm is, e.g., often used to solve linearized dynamic stochastic general equilibrium models, the workhorses of modern macroeconomics (King and Rebelo, 1999; Schmitt-Grohé and Uribe, 2004; Kollmann, 2015). Google Scholar records 2389 cites (09/2016) for the BK paper.

<sup>2</sup>Most likely this error is a typing or printing error. BK is a very classical paper, so a correct version of all key results in it should be available. BK's *general* routine for models with arbitrary numbers of states and jump variables (widely used in computer simulation packages) is accurate.

## 2 A LINEAR DYNAMIC MODEL WITH ONE STATE VARIABLE AND ONE JUMP VARIABLE

Consider the following model (the notation follows BK):

$$\begin{bmatrix} x_{t+1} \\ E_t p_{t+1} \end{bmatrix} = A \begin{bmatrix} x_t \\ p_t \end{bmatrix} + \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix} Z_t, \quad (1)$$

where  $x_t$  is a predetermined variable ('state'), and  $p_t$  is a non-predetermined ('jump') variable.  $Z_t$  is a  $(k \times 1)$  vector of exogenous variables.  $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$  is a  $(2 \times 2)$  matrix, and  $\gamma_1, \gamma_2$  are  $(1 \times k)$  vectors. Let  $\lambda_1, \lambda_2$  be the eigenvalues of  $A$ , with  $|\lambda_1| \leq |\lambda_2|$ , and let  $B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{bmatrix}$  be the matrix of eigenvectors of  $A$ , i.e.  $AB = BJ$ , with  $J = \begin{bmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{bmatrix}$ . Let  $C \equiv B^{-1}$ ,  $C = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \end{bmatrix}$ . Note that  $A = BJC$ . Proposition 1 of BK (p.1308) shows that model (1) has a unique (non-exploding) solution if and only if  $|\lambda_1| \leq 1$ ,  $|\lambda_2| > 1$ . BK (p.1309) state that then the solution of (1) is:

$$x_t = \lambda_1 x_{t-1} + \gamma_1 Z_{t-1} + \mu \sum_{i=0}^{\infty} \lambda_2^{-i-1} E_{t-1} Z_{t+i-1}, \quad (2)$$

$$p_t = a_{12}^{-1} \left[ (\lambda_1 - a_{11}) x_t + \mu \sum_{i=0}^{\infty} \lambda_2^{-i-1} E_t Z_{t+i} \right], \quad (3)$$

$$\text{with } \mu \equiv (\lambda_1 - a_{11}) \lambda_1 - a_{12} \lambda_2. \quad (4)$$

*Comment:* When  $\mu$  is given by (4), then  $\mu \sum_{i=0}^{\infty} \lambda_2^{-i-1} E_{t-1} Z_{t+i-1}$  and  $\mu \sum_{i=0}^{\infty} \lambda_2^{-i-1} E_t Z_{t+i}$  are  $(k \times 1)$  vectors; thus (2), (3) cannot hold for  $k > 1$  as  $x_t, p_t$  are scalars. As shown below BK's formula (4) for  $\mu$  is incorrect. The correct formula is:

$$\mu = (\lambda_1 - a_{11}) \gamma_1 - a_{12} \gamma_2. \quad (5)$$

*Proof:* (2) and (3) are special cases of the general solution for linear difference models (with arbitrary numbers of variables) in BK's Proposition 1. (See Appendix.) The general solution for state  $x_t$  shows that the correct expression for  $\mu$  in (2) is  $\mu = -(b_{11} \lambda_1 c_{12} + b_{12} \lambda_2 c_{22}) c_{22}^{-1} (c_{21} \gamma_1 + c_{22} \gamma_2)$ . Write this as  $\mu = \phi_1 \gamma_1 + \phi_2 \gamma_2$ , with  $\phi_1 \equiv -(b_{11} \lambda_1 c_{12} c_{22}^{-1} c_{21} + b_{12} \lambda_2 c_{21})$ ,  $\phi_2 \equiv -(b_{11} \lambda_1 c_{12} + b_{12} \lambda_2 c_{22})$ .  $A = BJC$  implies  $a_{11} = b_{11} \lambda_1 c_{11} + b_{12} \lambda_2 c_{21}$  and  $a_{12} = b_{11} \lambda_1 c_{12} + b_{12} \lambda_2 c_{22}$ . Thus  $\phi_2 = -a_{12}$ . Substituting  $b_{12} \lambda_2 c_{21} = a_{11} - b_{11} \lambda_1 c_{11}$  into the definition of  $\phi_1$  gives  $\phi_1 \equiv -(a_{11} + b_{11} \lambda_1 [c_{12} c_{22}^{-1} c_{21} - c_{11}])$ .  $B = C^{-1}$  implies  $b_{11} = c_{22} / (c_{11} c_{22} - c_{12} c_{21})$  and  $c_{12} c_{22}^{-1} c_{21} - c_{11} = -b_{11}^{-1}$ . Thus

$\phi_1 = \lambda_1 - a_{11}$ . This implies (5). The general solution for jump variable  $p_t$  shows that equation 3 holds when  $\mu$  is defined by (5). ■

## APPENDIX

## BLANCHARD AND KAHN (1980): THE GENERAL MODEL

Consider the model

$$\begin{bmatrix} X_{t+1} \\ E_t P_{t+1} \end{bmatrix} = A \begin{bmatrix} X_t \\ P_t \end{bmatrix} + \gamma Z_t, \quad (\text{A1})$$

where  $X_t$  is an  $n \times 1$  vector of predetermined variable, and  $p_t$  is an  $m \times 1$  vector of non-predetermined variables;  $Z_t$  is a  $(k \times 1)$  vector of exogenous variables.  $A$  is an  $(n+m) \times (n+m)$  matrix, and  $\gamma$  is an  $(n+m) \times k$  matrix. Consider the Jordan canonical form  $A = C^{-1}JC$ , where  $C$  and  $J$  are  $(n+m) \times (n+m)$  matrices. Let the diagonal elements of  $J$  (i.e. the eigenvalues of  $A$ ) be ordered by increasing absolute value. Let  $\bar{n}$  ( $\bar{m}$ ) denote the number of eigenvalues of  $A$  that are on or inside the unit circle (outside the unit circle). Partition  $J$  as  $J = \begin{bmatrix} J_1 & 0 \\ 0 & J_2 \end{bmatrix}$ , where  $J_1$  and  $J_2$  are matrices of dimensions  $(\bar{n} \times \bar{n})$  and  $(\bar{m} \times \bar{m})$  respectively. Decompose  $C$ ,  $B \equiv C^{-1}$  and  $\gamma$  as  $C = \begin{bmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{bmatrix}$ ,  $B = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$  and  $\gamma = \begin{bmatrix} \gamma_1 \\ \gamma_2 \end{bmatrix}$ , where  $C_{11}, C_{12}, C_{21}, C_{22}$  are matrices of dimensions  $(\bar{n} \times n)$ ,  $(\bar{n} \times m)$ ,  $(\bar{m} \times n)$  and  $(\bar{m} \times m)$  respectively;  $B_{11}, B_{12}, B_{21}, B_{22}$  have dimensions  $(n \times \bar{n})$ ,  $(n \times \bar{m})$ ,  $(m \times \bar{n})$  and  $(m \times \bar{m})$  respectively, while  $\gamma_1$  and  $\gamma_2$  have dimensions  $(\bar{n} \times k)$  and  $(\bar{m} \times k)$  respectively. Proposition 1 in Blanchard and Kahn (1980) states that the model (A1) has a unique (non-explosive) solution if and only if the number of non-predetermined variables equals the number of eigenvalues of  $A$  outside the unit circle:  $m = \bar{m}$ . If that condition is met, then the solution is:

$$\begin{aligned} X_t &= B_{11} J_1 B_{11}^{-1} X_{t-1} + \gamma_1 Z_{t-1} - (B_{11} J_1 C_{12} + B_{12} J_2 C_{22}) \\ &\quad \times C_{22}^{-1} \sum_{i=0}^{\infty} J_2^{-i-1} (C_{21} \gamma_1 + C_{22} \gamma_2) E_{t-1} Z_{t+i-1}, \\ P_t &= -C_{22}^{-1} C_{21} X_t - C_{22}^{-1} \sum_{i=0}^{\infty} J_2^{-i-1} (C_{21} \gamma_1 + C_{22} \gamma_2) E_t Z_{t+i}. \end{aligned}$$

## REFERENCES

- Blanchard, O. and Kahn, C. (1980). 'The Solution of Linear Difference Models Under Rational Expectations', *Econometrica*, Vol. 48, pp. 1305–1311.
- King, R. and Rebelo, S. S. (1999). 'Resuscitating Real Business Cycles', in J. Taylor and M. Woodford (eds), *Handbook of Macroeconomics*, Vol. 1B, Elsevier: Amsterdam, pp. 927–1007.
- Kollmann, R. (2015). 'Tractable Latent State Filtering for Non-Linear DSGE Models Using a Second-Order Approximation and Pruning', *Computational Economics*, Vol. 45, pp. 239–260.
- Schmitt-Grohé, S. and Uribe, M. (2004). 'Solving Dynamic General Equilibrium Models Using A Second-Order Approximation to the Policy Function', *Journal of Economic Dynamics and Control*, Vol. 28, pp. 755–775.